

EXHIBIT F

Health Consultation

EXPOSURE INVESTIGATION

HERCULANEUM LEAD SMELTER SITE
(A/K/A DOE RUN LEAD SMELTER)

CITY OF HERCULANEUM, JEFFERSON COUNTY, MISSOURI

EPA FACILITY ID: MOD006266373

JUNE 9, 2005

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
Agency for Toxic Substances and Disease Registry
Division of Health Assessment and Consultation
Atlanta, Georgia 30333

Health Consultation: A Note of Explanation

An ATSDR health consultation is a verbal or written response from ATSDR to a specific request for information about health risks related to a specific site, a chemical release, or the presence of hazardous material. In order to prevent or mitigate exposures, a consultation may lead to specific actions, such as restricting use of or replacing water supplies; intensifying environmental sampling; restricting site access; or removing the contaminated material.

In addition, consultations may recommend additional public health actions, such as conducting health surveillance activities to evaluate exposure or trends in adverse health outcomes; conducting biological indicators of exposure studies to assess exposure; and providing health education for health care providers and community members. This concludes the health consultation process for this site, unless additional information is obtained by ATSDR which, in the Agency's opinion, indicates a need to revise or append the conclusions previously issued.

You May Contact ATSDR TOLL FREE at
1-888-42ATSDR

or

Visit our Home Page at: <http://www.atsdr.cdc.gov>

Table of Contents

| | |
|--------------------------------|----|
| <i>Executive Summary</i> | 4 |
| <i>Target Population</i> | 6 |
| <i>Test Procedures</i> | 7 |
| <i>Biological Samples</i> | 8 |
| <i>Environmental Samples</i> | 9 |
| <i>Results and Discussion</i> | 9 |
| <i>Biological Samples</i> | 9 |
| <i>Cadmium</i> | 9 |
| <i>Arsenic</i> | 11 |
| <i>Total Urine Arsenic</i> | 12 |
| <i>Inorganic Urine Arsenic</i> | 13 |
| <i>Environmental Samples</i> | 14 |
| <i>Fruits and Vegetables</i> | 14 |
| <i>Arsenic</i> | 14 |
| <i>Cadmium</i> | 14 |
| <i>Lead</i> | 15 |
| <i>Reporting Results</i> | 17 |
| <i>Conclusions</i> | 18 |
| <i>Recommendations</i> | 18 |
| <i>Authors</i> | 20 |
| <i>References</i> | 22 |

Abbreviations and Acronyms

| | |
|-----------|---|
| ATSDR | Agency for Toxic Substances and Disease Registry |
| ACGIH | American Conference of Governmental Industrial Hygienists |
| CDC | Centers for Disease Control and Prevention |
| EI | exposure investigation |
| EPA | U.S. Environmental Protection Agency |
| FDA | Food and Drug Administration |
| G | grams |
| MDHSS | Missouri Department of Health and Senior Services |
| NCEH | National Center for Environmental Health (CDC) |
| mg/kg/day | milligrams per kilogram (of body weight) per day |
| mL | milliliter |
| MRL | minimal risk level |
| ppm | parts per million |
| PTTIL | Provisional Total Tolerable Intake Level |
| USDA | U.S. Department of Agriculture |
| µg/L | micrograms per liter |

vegetables were preferentially selected for testing because they tend to take up more cadmium (ATSDR 1999b). The risk posed by eating cadmium-containing produce would depend on the rate of consumption. For example, an adult (70 kg) might eat 1 cup of lettuce (56 grams) per day, the vegetable with the highest concentration of cadmium (US Dept. of Agriculture Database 2004).

Using the highest cadmium concentration of 0.391 ppm, which was detected in lettuce, the maximum cadmium ingestion rate for this scenario would be:

$$0.391 \mu\text{g/g (ppm)} \times 56 \text{ grams} \div 70 \text{ kg body weight} = 0.3128 \mu\text{g/kg/day}$$

The result, 0.3128 $\mu\text{g/kg/day}$, is equivalent to 0.00031 mg/kg/day.

This estimated maximum cadmium dose slightly exceeds the ATSDR chronic MRL for cadmium of 0.0002 mg/kg/day. Exposure to a level above the MRL does not mean that adverse health effects will occur. MRLs serve as a screening tool to help public health professionals decide where to look more closely for potential health risks. An MRL is an estimate of the daily human exposure to a hazardous substance that is likely to be without appreciable risk of adverse noncancer health effects over a specified duration of exposure.

The ATSDR chronic MRL for cadmium is based on exposure for 365 days a year for many years. However, the participants in this investigation ate home-grown vegetables occasionally, during a few months of the year, so their average daily dose would be considerably less than the value calculated above. Because of the large surface area of lettuce, a portion of the cadmium concentration may result from surface deposition. Peeling and thoroughly washing vegetables would reduce the amount of surface metal contamination.

The consumption of these fruits and vegetables is not likely to result in adverse health problems from cadmium exposure if consumed occasionally for a few months each year. To be protective of public health, ATSDR recommends that individuals follow general food safety guidelines. Those include washing hands before preparing foods and washing foods before consumption. Peeling and thoroughly washing vegetables would reduce the amount of surface metal contamination.

Lead

Many plants can take up lead from the soil (ATSDR 1999a). Edible plants acquire lead from the soil through their roots, by direct foliage uptake, and by surface deposition of particulate matter (ATSDR 1999a). Lead generally deposits superficially on leaves, with less leaf penetration than cadmium (Vousta 1996). At significantly high levels, lead is poisonous to plants and can stunt plant growth.

The lead levels in vegetables and fruits tested ranged from 0.008 – 1.737 ppm. Green leafy vegetables tend to accumulate lead on the plant surface. Root vegetables take up

lead into the plant tissue (ATSDR 1999a; Finster et al. 2003). The risk posed by eating lead-containing produce depends on the frequency and amount of consumption. An ATSDR MRL for lead is not available. Therefore, the U.S. Food and Drug Administration (FDA) provisional tolerable lead intake levels were used.

The FDA's recommended Provisional Total Tolerable Intake Level (PTTIL) for lead in children less than 6 years of age is 6 μg lead/day. For children 7 years and older, the PTTIL is 15 μg lead/day. It increases to 75 μg lead/day for adults (USFDA 1993).

The maximum concentration of lead detected in a plant in this investigation—1.737 ppm—was detected in lettuce. Plants with higher surface areas (green leafy vegetables, such as lettuce, collard greens, polk wild greens and swiss chard) tend to have higher lead levels (Finster et al. 2003). Contaminated soil and dust attaches to the plant surface more easily and tends to remain on vegetables that are not washed well.

The risk posed by eating lead-containing produce would depend on the rate of consumption. For instance, an adult might eat 1 cup of lettuce (56 grams) per day (USDA 2004). Using the maximum lead level detected (1.737 ppm), an adult eating lettuce at that rate would ingest about 95 μg lead/day, which exceeds the adult PTTIL for lead. This ingestion estimate also exceeds the lead PTTIL for children. The other vegetables contained less than half the lead concentration found in lettuce. Most of the individuals who provided the home-grown fruits and vegetables only ate them when ripe and in season. This would result in a few months of occasional consumption each year.

Based on these assumptions, consumption of home-grown green leafy vegetables should be limited to a few times a week when in season. Lead content in lettuce, which has a large surface area, may partly result from deposition of lead-contaminated soil on the leaf surfaces.

To be protective of public health, ATSDR recommends that adults limit their consumption of green leafy vegetables to a few times a week when in season. Because young children and pregnant women are more sensitive to the effects of lead, they should avoid eating home-grown leafy vegetables. To reduce potential exposures to lead, home gardeners should preferentially plant fruiting vegetables instead of green leafy vegetables. Peeling and thoroughly washing vegetables can reduce the amount of lead contamination.

Reporting Results

In April 2004, individual test results and an explanation of their significance were provided to the participants of this investigation. An ATSDR physician was available to discuss participants' results.

Reviewed by:

Susan Metcalf, M.D., M.P.H.
Team Leader
Exposure Investigations and Consultations Branch
Division of Health Assessment and Consultation

Susan Moore, M.S.
Acting Branch Chief
Exposure Investigations and Consultations Branch
Division of Health Assessment and Consultation

Acknowledgments

Michael Lewin—ATSDR, Division of Health Studies, for statistical analysis.

Duane Hughes—Food and Drug Administration, for vegetable processing and analysis.

Ken Orloff, Ph.D. and John Wheeler, Ph.D.—ATSDR, Division of Health Assessment and Consultation, for technical assistance with vegetable analysis.

Lynn Wilder, CIH—ATSDR, Division of Health Studies, for technical guidance on environmental sampling.

Kathleen Caldwell, Ph.D. and Robert Jones, Ph.D.—Centers for Disease Control and Prevention, for urine cadmium and arsenic laboratory analysis.

Kim Elmore, Ph.D.—ATSDR, for geographical information systems.

References

Agency for Toxic Substance and Disease Registry (ATSDR). 1999a. Toxicological profile for lead (update). Atlanta: US Department of Health and Human Services; 1999 Jul.

Agency for Toxic Substance and Disease Registry (ATSDR). 1999b. Toxicological profile for cadmium (update). Atlanta: US Department of Health and Human Services; 1999 Jul.

Agency for Toxic Substance and Disease Registry (ATSDR). 2000. Toxicological profile for arsenic (Update). Atlanta: US Department of Health and Human Services; 2000 Sep.

Agency for Toxic Substances and Disease Registry (ATSDR). 2002a. Health consultation on blood lead results for 2001 calendar year, Herculaneum lead smelter site, Herculaneum, Missouri. Atlanta: US Department of Health and Human Services; 2002 Feb 26.

Agency for Toxic Substances and Disease Registry (ATSDR). 2002b. Health consultation on determination if remedial actions are protective of public health, Herculaneum lead smelter site, Herculaneum, Missouri. Atlanta: US Department of Health and Human Services; 2002 Apr 16.

Agency for Toxic Substances and Disease Registry (ATSDR). 2002c. Health consultation on arsenic and cadmium levels in air and residential soils, Herculaneum Lead Smelter Site, Herculaneum, Missouri. Atlanta: US Department of Health and Human Services; 2002 Nov 12.

American College of Industrial Hygienists (ACGIH). 2001. Cadmium and inorganic compounds. In: Documentation of the biological exposure indices. 7th ed. Cincinnati: ACGIH Worldwide. p. 1–10.

Andren P, Schutz A., Vahter M., Attewell R., Johansson L., Willers S., Skerfving S. 1988. Environmental exposure to lead and arsenic among children living near a glassworks. *Science of the Total Environment*. 77(1):25-34.

Buchet JP, Lauwerys R, Roels H, Bernard A, Brauaux P, Claeys F, et al. 1990. Renal effects of cadmium body burden in the general population. *Lancet* 336:699–702.

Centers for Disease Control and Prevention. 2003 Jan 31. National Center for Environmental Health: Second National Report on Human Exposure to Environmental Chemicals. Available at URL: <http://www.cdc.gov/exposurereport/pdf/SecondNER.pdf>. Accessed April 2005.

- Finster ME, Gray KA, Binns HJ. 2004. Lead levels of edibles grown in contaminated residential soils: a field survey. *Sci Total Environ.* 320(2-3):245–57.
- Hotz P, Buchet JP, Bernard A, Lison D, Lauwerys R. 1999. Renal effects of low-level environmental cadmium exposure: 5-year follow-up of a subcohort from the Cadmibel study. *Lancet.* 354:1508–13.
- Jarup L, Bergland M, Elidner CG, Nordberg G, Vahter M. 1998. Health effects of cadmium exposure: a review of the literature and a risk estimate. *Scand J Work Environ Health* 24(suppl 1):7-51.
- Jensen GE, Christensen JM., Poulsen OM. 1991. Occupational and environmental exposure to arsenic--increased urinary arsenic level in children. *Science of the Total Environment.* 107:169-77.
- Johnson LR and Farmer JG. 1989. Urinary arsenic concentrations and speciation in Cornwall residents. *Environ Geochem Health* 11:39-44.
- Kallman, DA, Hughes J, Van Belle G, Burbacher T, Bolgiano D, Coble K, et al. 1990. The effect of variable environmental arsenic contamination on urinary concentrations of arsenic species. *Environ Health Perspect* 89:145–51.
- Lauwerys, RR, Bernard A, Roels H, Buchet JP. 1994. Cadmium: exposure markers as predictors of nephrotoxic effects. *Clin Chem* 40(7):1391–4.
- Missouri Department of Natural Resources (MDNR). 1999 Mar 30. Preliminary assessment: Herculanum lead smelter site, Jefferson County, Missouri. Jefferson City, Missouri: Missouri Department of Natural Resources.
- Paschal DC, Burt V, Caudill SP, Gunter EW, Pirkle JL, Sampson EJ, et al. 2000. Exposure of the US population aged 6 years and older to cadmium:1988–1994. *Arch Environ Contam Toxicol* 38:377–83.
- Sullivan JB and Krieger GR. 2001. Clinical environmental health and toxic exposures. 2nd ed. Philadelphia: Lippincott Williams & Williams. Page 862.
- Trevesian A, Nicoletto G, Maso S, Grandesso G, Odyne A, Secondin L. 1994. Biological monitoring of cadmium exposure: reliability of spot urine samples. *Int Archives Occup Environ Health* 65:373–5.
- US Department of Agriculture, Agricultural Research Service. 2004. USDA National Nutrient Database for Standard Reference, Release 16-1; Nutrient Data Laboratory. Available at URL: <http://www.nal.usda.gov/fnic/foodcomp>. Accessed April 2005.

US Food and Drug Administration. 1993. Guidance document for lead in shellfish. Washington, DC: Center for Food Safety and Applied Nutrition. Available at URL: <http://www.cfsan.fda.gov/~frf/guid-pb.html#sVI>.

Vousta D, Grimanis A, Samara C. 1996. Trace elements in vegetables grown in an industrial area in relation to soil and air particulate matter. *Environ Pollution* 94(3):325–35.

Wittman R and Hu, H. 2002. Cadmium exposure and nephropathy in a 28-year-old female metals worker. *Environ Health Perspect* 110(12):1261–6.

EXHIBIT G

EDMUND G. BROWN JR.
Attorney General

State of California
DEPARTMENT OF JUSTICE



1515 CLAY STREET, 20TH FLOOR
P.O. BOX 70550
OAKLAND, CA 94612-0550

Public: (510) 622-2100
Telephone: (510) 622-2149
Facsimile: (510) 622-2270
E-Mail: Ed.Weil@doj.ca.gov

March 3, 2008

JL Sean Slattery
Del Mar Law Group, LLP
322 8th Street, Suite 101
Del Mar, CA 92014

David Lavine
Laralei Paras
Hirst & Chanler
2560 Ninth Street
Parker Plaza, Suite 214
Berkeley, CA 94710

RE: Proposition 65 Claims Concerning Lead in Lipstick

Dear Mr. Slattery, Mr. Lavine, and Ms. Paras:

The release of a report titled "A Poison Kiss: The Problem of Lead in Lipstick," by the Campaign for Safe Cosmetics in October of 2007 ("CSC Report"), has raised public concern about whether the presence of lead in lipstick violates Proposition 65. On January 22, 2008, Christine Deubler, represented by Del Mar Law Group, served notices of violation under Proposition 65 alleging that L'Oreal USA, Maybelline, LLC, Parfums Christian Dior, Procter & Gamble, Inc., and Peacekeeper Cause-Metics, have violated Proposition 65 by failing to warn of the presence of lead in certain lipsticks. In the latter half of 2007, the Attorney General also received approximately ten notices of violation from Whitney Leeman, represented by Hirst & Chanler, alleging that various companies have failed to warn concerning lead in various cosmetics, some of which are lipsticks. Some of those notices pre-date the CSC Report, but raise many similar issues. Ms. Leeman has served additional notices raising similar claims as recently as February 19, 2008.

Over the past few weeks, we have exchanged information concerning the merits of these claims with the law firms representing the noticing parties. Del Mar Law Group has today advised us that they will not proceed with claims raised in their notices, which we consider an appropriate and responsible decision. Hirst & Chanler has indicated that they intend to continue pursuing these claims.

Mr. Slattery, Mr. Lavine, Ms. Paras

March 3, 2008

Page 2

As you may know, the Attorney General has brought numerous actions enforcing Proposition 65 with respect to lead in consumer products, including imported candies, soft drink bottles, toys, jewelry, tableware, drinking water faucets, and calcium supplements. Thus, he is committed to protecting California consumers from products containing lead in excess of legal requirements, and vigorously enforcing Proposition 65.

Ordinarily, when the Attorney General declines to pursue a proposed private Proposition 65 enforcement matter, it should not be considered a rejection of the claim on the merits. Nor does the Attorney General typically make a public statement concerning the merits of the matter. In a few instances, however, where an issue is of great concern to the public, it is appropriate to explain that we have concluded that no action is warranted based on the facts of the case. We believe that this is one of those instances.

Accordingly, this letter sets forth why, after thorough analysis, we have concluded that lead in lipstick at the levels identified in the CSC Report, and up to 5 parts per million lead, does not raise a reasonable claim of a Proposition 65 violation and ought not to be pursued.

A. Analysis

1. Lead Contents.

The CSC report tested 33 samples of lipstick and found 13 with no detectable lead (at a detection limit of 0.02 ppm), 9 with less than 0.1 ppm, and 11 within a range of more than 0.1 to as much as 0.65. This data is consistent with what has been found on previous occasions. In other instances, higher levels, in the 1-3 ppm range, have been reported.

2. Proposition 65 Requirements.

Proposition 65 requires that a business that "knowingly and intentionally" exposes persons to chemicals known to the state to cause cancer or reproductive harm must provide a "clear and reasonable warning," unless the business can prove that the exposure would have "no observable effect assuming exposure at one thousand (1,000) times the level in question," for reproductive toxicants, or pose "no significant risk" for carcinogens. (Health & Saf. Code, §§ 25249.6, 25249.10, subd. (c).) Designation of chemicals as known to cause cancer or reproductive toxicity, and regulatory standards for compliance, are implemented by the designated lead agency, the Office of Environmental Health Hazard Assessment ("OEHHA").

Lead is listed as a reproductive toxicant and a carcinogen under Proposition 65. (Cal. Code Regs., tit. 22, § 12000, subd. (a), (b).) For a number of reproductive toxicants, OEHHA

Mr. Slattery, Mr. Lavine, Ms. Paras
 March 3, 2008
 Page 3

has adopted a "safe-harbor" Maximum Allowable Dose Level (MADL). Where such a level is adopted, "[e]xposure to a chemical at a level which does not exceed the level set forth in [the regulation] for such chemical has no observable effect assuming exposure at one thousand (1,000) times that level." (Cal. Code Regs., tit. 22, § 12805, subd. (a).) In other words, if the exposure is less than the safe-harbor level, no warning is required. If the exposure exceeds the safe-harbor level, the business still has the option of attempting to prove a different (presumably higher) Maximum Allowable Dose Level, but must do so in compliance with scientific standards set forth in the regulations. (Cal. Code Regs., tit. 22, § 12801.) The adopted safe-harbor MADL for lead is 0.5 micrograms per day. (Cal. Code Regs., tit. 22, § 12805, subd. (b).)¹

This daily dose does **not** directly correspond to the concentration of lead in the product, however. To determine whether a warning is required, one must analyze whether the "reasonably anticipated rate of intake or exposure for average users of the consumer product" would be less than 0.5 micrograms per day. (Cal. Code Regs., tit. 22, § 12821, subd. (d)(2).) This requires a determination of how much lipstick people use.²

The meaning of "average users" is not fully settled. The only reported case on the issue upheld a trial court decision finding no violation where the 75th-85th percentile of users were below the warning level as an acceptable definition of average users. (*DiPirro v. Bondo Corporation* (2007) 153 Cal.App.4th 150, 190.) In other instances, courts have focused on the mean or the median.

3. Exposure Analysis.

a. Available Evidence.

There is some information available concerning average users of lipstick in the peer-

¹We think it is unlikely that a defendant could successfully prove a higher MADL for lead. In addition, although lead also is listed as a carcinogen, it is far less potent as a carcinogen than as a reproductive toxin, with a safe-harbor No Significant Risk Level of 15 micrograms per day. (Cal. Code Regs., tit. 22, § 12705, subd. (b)(1).) Thus, if the product does not require a warning for reproductive toxicity, it clearly will not require one for cancer.

²The CSC report suggested that a concentration of lead of 0.1 ppm is an appropriate standard under Proposition 65. That standard was borrowed from another case brought by the Attorney General concerning candy contaminated with lead, and was based on a variety of factors, most importantly concerning the amount of candy ingested by children. Since people do not eat lipstick the way children eat candy, the two analyses of exposure are quite different.

Mr. Slattery, Mr. Lavine, Ms. Paras
 March 3, 2008
 Page 4

reviewed literature. The first is a report prepared in 2001 by the Cosmetic, Toiletry, and Fragrance Association ("CTFA"). This data was admitted into evidence in 2004 in a trial of a private Proposition 65 enforcement case in which the trial court found that lipstick did not require a warning. (*DiPirro v. J.C. Penney*, San Francisco Superior Ct. No. 407150 [Statement of Decision filed February 9, 2005, pp. 37-47, 113-117].)³ (The products at issue in that case included lipstick, make-up, and leaded glassware, and the private plaintiff prevailed on the glassware issue.) The study subsequently was published in peer-reviewed literature. ("Exposure data for cosmetic products: lipstick, body lotion, and face cream," *Food and Chemical Toxicology* v. 43, pp. 279-291 (2005).)

The study estimated exposure by getting 311 women to participate, giving them lipstick, having them come back in two weeks, asking how many times per day they used it, and actually measuring the difference in the net weight of the lipstick. (Measuring the amount actually used has a significant advantage over simply asking the respondents how much they use, because it is not subject to memory error.) The study was reasonably well done given the inherent limitations involved, although there are some valid criticisms (e.g., some of the respondents apparently did not like the product, and therefore may not have used as much of it as they ordinarily would).

The study found that the mean number of applications was 2.4 times per day, with 11% of users applying it 4 or more times per day, and a median of 2.1 times per day. The mean amount of lipstick applied per application was 5 milligrams, with 12% of users applying 20 milligrams or more per application. Thus, using the means, the daily use of lipstick is 24 milligrams. Someone in both the top 11% of frequency of use and the top 12% of amount per application would use 80 milligrams each day.

Recently, two more studies were published in peer-reviewed literature, based on surveys conducted in Europe.⁴ These studies combined some large existing European databases in which thousands of persons kept diaries and answered questionnaires concerning cosmetic use, in combination with another study in which the actual use of lipstick in milligrams is measured. In this study, the arithmetic mean of use was 24.6 mg/day, the median was 17.1, the 80th percentile

³The plaintiff appealed, but the matter was settled while the appeal was pending.

⁴The two studies are "Probabilistic modeling of European consumer exposure to cosmetic products," *Food and Chemical Toxicology* 45 (2007) 2086-2096, and "European consumer exposure to cosmetic products, a framework for conducting population exposure assessments," *Food and Chemical Toxicology* 45 (2007) 2097-2108.

Mr. Slattery, Mr. Lavine, Ms. Paras
 March 3, 2008
 Page 5

was 39.7, and the 90th percentile was 56.5 mg/day.⁵

In the *DiPirro v. J.C. Penney* case referenced above, the plaintiff relied on an expert who calculated exposures based on an average number of applications per day of six times, taken from a figure presented in the U.S. EPA's Exposure Factors Handbook, and estimated the amount of lipstick applied through an estimate of the surface area of the lips and the depth of the application at 53 milligrams per application. This results in exposure of 318 milligrams per day. The court in that case rejected the expert's testimony, mostly for reasons not directly related to this aspect of her opinions. In any event, the expert's method of estimating offered in that case does not seem to be supported by any other data.

While not a matter of formal evidence, it is worth considering whether these exposure parameters seem to fit with ordinary understanding of the consumption of the products. An average tube of lipstick weighs about 0.14 ounces (about 4 grams, or 4,000 milligrams). At 24 milligrams a day, it would take about 5 ½ months to use a tube, if that is the only tube used. If we assume the use is 100 milligrams per day, it would take 40 days to use a tube, if that is the only tube used. If one assumes the use is 300 milligrams per day, a tube would be consumed in less than two weeks. On that basis, consumption estimates of up to 100 milligrams per day seem consistent with common experience.

b. Likely Exposure Estimate.

The above-referenced data suggests that the highest use supportable by the facts and the regulation would be about 100 milligrams per day. (Indeed, this figure exceeds the 90th percentile as indicated in any of the published studies.) Comparing this to the 0.5 microgram per day Safe-harbor level, this means that the lipstick would need to contain 5.0 ppm lead to require a warning.

The analysis above estimates the total amount of lead the user comes into contact with, i.e., gets on the lips. Under the statute and regulations, any contact with the body constitutes an "exposure," which is sufficient to make the prima facie case. (Cal. Code Regs., tit. 22, § 12102, subd. (i); *Consumer Cause v. Weider Nutrition* (2001) 92 Cal.App.4th 363.) Attempting to prove under the statute that the exposure has "no observable effect . . . at the level in question," a

⁵These studies computed "daily averages" of use. For a reproductive toxicant such as lead, the usage on a given day is considered the appropriate measure of exposure, not the long-term daily average. In these studies, however, the product was used every day, and there was no evidence that the usage varied significantly from day-to-day. Thus, in this case there is little practical difference in the two numbers.

Mr. Slattery, Mr. Lavine, Ms. Paras
 March 3, 2008
 Page 6

defendant can attempt to prove that the substance in question is not actually absorbed by the body, which would affect the level at which the warning exemption would apply. In this instance, defendants may try to prove that in fact much of the lipstick is never actually absorbed by the body, because relatively little of it is actually absorbed through the lips (lead is not well-absorbed through skin), or swallowed and ingested. We have seen no reliable data on how much lipstick is actually ingested, although reason suggests that it is some fraction of the total amount applied. Thus, if the matter were to proceed to litigation, it seems likely that some allowance would have to be made for the fact that not all of the product is ingested or absorbed. In this analysis, we have not eased the standard to account for any claim that lead is not absorbed, which effectively applies a stricter and more health-protective standard than might be applied by a court after trial.⁶

B. Conclusion

The above analysis indicates that lipstick with lead concentrations at the levels found in the CSC report could not plausibly be considered to trigger a duty to warn under Proposition 65. Indeed, it appears that a reasonable claim that there is a duty to warn would not arise until concentrations reached 5 ppm lead. We have not seen any publicly-available, reliable data showing significant amounts of lipstick with lead concentrations above those levels. Accordingly, based on the data we have reviewed, we do not think Proposition 65 actions would be warranted for lipstick with lead levels less than 5 ppm. If data showing higher concentrations of lead in lipstick are provided, the matter would need to be evaluated in the light of that information.

In addition to the public concern about the matter, we are concerned about the potential

⁶A defendant seeking to raise this issue may not be able to rely on the 0.5 microgram per day MADL, however. The regulations allow a defendant to establish the "no observable effect" defense in one of two ways, either "(1) By means of an assessment that meets the standards described in Section 12803" or "(2) By application of a specific regulatory level for the chemical in question as provided in Section 12805." (Cal. Code Regs., tit. 22, § 12801, subd. (b).) The safe-harbor MADLs provided in section 12805 by their terms apply to the amount of "exposure" to the chemical (Cal. Code Regs., tit. 22, § 12805, subd. (a)), which in this instance is the amount of lipstick applied. If the exposure exceeds the safe-harbor level, but the defendant argues that there will be no actual effect because of a lack of absorption or ingestion, the defendant is no longer proceeding under safe-harbor provision, but must establish the No Observable Effect defense under section 12801(b)(1), by means of an independent assessment. Of course, the evidence that led OEHHA to establish the safe-harbor MADL would be considered as part of the defense, and, depending on the facts, could lead to the same result.

Mr. Slattery, Mr. Lavine, Ms. Paras
March 3, 2008
Page 7

use of this claim by plaintiffs to pursue matters in a manner that does not promote the public interest. (*Consumer Defense Group v. Rental Housing Industry Members* (2006) 137 Cal.App.4th 1185.) Lipstick is sold in literally thousands of stores throughout the state. In the past, some plaintiff groups have pursued Proposition 65 litigation against large numbers of small businesses, with little evidence of whether the retailer has knowledge of the presence of the listed chemical in the product or on the premises. If this were to occur here, many of those stores would find it more practical to pay a small settlement to the plaintiff than to contest the case. Those stores also might post warnings for products that clearly do not require warnings, which is not in the public interest. Nor would such proceedings result in the "enforcement of an important right affecting the public interest" or confer a "significant benefit" on the general public as those terms are used in Code of Civil Procedure section 1021.5.

We hope this objective review of the merits of the issue will discourage your client and any other private plaintiffs from pursuing these matters. If such cases are brought, we will carefully review them in order to determine whether it is appropriate for the Attorney General to take any action.

Sincerely,

A handwritten signature in black ink, appearing to read "E. G. Weil", written in a cursive style.

EDWARD G. WEIL
Supervising Deputy Attorney General

For EDMUND G. BROWN JR.
Attorney General

cc: Linda Katz, M.D. (Director, Office of Cosmetics and Colors, FDA/CFSAN)
Elizabeth Anderson, Personal Care Products Council